

View this article online at: patient.info/doctor/six-item-cognitive-impairment-test-6cit

## Six-item Cognitive Impairment Test (6CIT)

The 6-item Cognitive Impairment Test (6CIT) Kingshill Version 2000® was developed in 1983. <sup>[1]</sup> This was by regression analysis of the Blessed Information Memory Concentration Scale (BIMC). <sup>[2]</sup> The 6CIT is a useful dementia screening tool in Primary Care. It was used in a large European assessment tool (Easycare©) and with new computerised versions its usage is increasing.

Six-item Cognitive Impairment Test (6CIT) - Kingshill Version 2000		
1. What year is it?	Correct - 0 points Incorrect - 4 points	
2. What month is it?	Correct - 0 points Incorrect - 3 points	
3. Give the patient an address phrase to remember with 5 components - eg: John, Smith, 42, High St, Bedford.		
4. About what time is it (within one hour)?	Correct - 0 points Incorrect - 3 points	
5. Count backwards from 20-1.	Correct - 0 points 1 error - 2 points More than one error - 4 points	
6. Say the months of the year in reverse.	Correct - 0 points 1 error - 2 points More than one error - 4 points	
7. Repeat address phrase.	Correct - 0 points 1 error - 2 points 2 errors - 4 points 3 errors - 6 points 4 errors - 8 points All wrong - 10 points	
6CIT score = /28		

Many thanks to Dr Patrick Brooke, General Practitioner & Research Assistant in Dementia for his help with the original article. The Kingshill Research Centre, Swindon, UK owns the copyright to The Kingshill Version 2000 of the 6CIT but allows free usage to healthcare professionals.

- Number of questions: 6.
- Time taken to perform: 3-4 minutes.
- **Score**: the 6CIT uses an inverse score and questions are weighted to produce a total out of 28. Scores of 0-7 are considered normal and 8 or more significant.
- **Advantages**: the test has high sensitivity without compromising specificity, even in mild dementia. It is easy to translate linguistically and culturally.

- **Disadvantages**: the main disadvantage is in the scoring and weighting of the test, which is initially confusing; however, computer models have simplified this greatly.
- **Probability statistics**: at the 7/8 cut-off: Overall figures sensitivity = 90%, specificity = 100%; in mild dementia, sensitivity = 78%, specificity = 100%.

The 6CIT is a much newer test than the Abbreviated Mental Test (AMT). It would appear to be culturally and linguistically translatable with good probability statistics; however, it is held back by its more complex scoring system. Furthermore, it would be nice to see some additional larger population studies using the test.

## Further reading & references

- 1. Brooke P, Bullock R; Validation of a 6 item cognitive impairment test with a view to primary care usage. Int J Geriatr Psychiatry, 1999 Nov;14(11):936-40.
- 2. Blessed G, Tomlinson BE, Roth M, The association between quantitative measures of dementia and of senile change in the cerebral grey matter of elderly subjects. Br J Psychiatry. 1968 Jul;114(512):797-811.

**Disclaimer:** This article is for information only and should not be used for the diagnosis or treatment of medical conditions. EMIS has used all reasonable care in compiling the information but makes no warranty as to its accuracy. Consult a doctor or other healthcare professional for diagnosis and treatment of medical conditions. For details see our conditions.

Original Author:	Current Version:	Peer Reviewer:
Dr Huw Thomas	Dr Hayley Willacy	Dr Colin Tidy
Document ID: 4172 (v8)	Last Checked: 28/01/2016	Next Review: 26/01/2021

View this article online at: patient.info/doctor/six-item-cognitive-impairment-test-6cit

Discuss Six-item Cognitive Impairment Test (6CIT) and find more trusted resources at Patient.

## Ask your doctor about Patient Access

Book appointments

Order repeat prescriptions

View your medical record

Create a personal health record (105 only)







Simple, quick and convenient.

Visit patient.info/patient-access
or search 'Patient Access'

© Patient Platform Limited - All rights reserved.